

h 44
h 69
h 74.75
89

96

1633

138
140

23 tints of plumage vary with climate

69 Billson on variation of Beak in

Tetrao saliceti.

(2)

70. many short-tailed birds have 1 or 2 pairs of
glo-canal. (as Kingfisher) can then

be compensation or rudimentary.
(allude when I talk of important organs
being few & not variable.)

69 References to Bonchi's papers. I have rec^d
2 Jan acc^y Bonch.

74 on changes of Ducks wings & feet, tamed

103 on speeding of sparrow with collection in

Peripia


143 Corvus of Faroe & C. cornix

Appendix systematic at end on doubtful species

157 on *Tetrao saliceti* & *scoticus*

being same species (2)

(249)



722. on variability of Head & Beak

Brehm made some of his species from single
specimens & by specimens.

722. Cymbium 18 or 20 tars feathers

731 Beak & length of ^{primaries} wings in Anser segetum
variable - so is it with Tormenter geese

733 tars feathers increase in number in
Cymbium with age

Is Anser segetum
lapping back of
Tormenter geese.

Bruch in Isis 1828 Band XXI

to Isis 1829 Band XXII

p. 629 Caudals increase with age. variable
in Anser segetum - 16 - 20!

p. 152 on the Sparrow Cl. 4

(249)

Handwritten notes, possibly a list or a set of instructions, written in a cursive script. The text is difficult to decipher due to the fading and the angle of the writing.

PAC